

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44 // (A61K 31/44, 31:215) (A61K 31/44, 31:19) (A61K 31/44, 31:35) (A61K 31/44, 31:22) (A61K 31/44, 31:44)		A1	(11) International Publication Number: WO 97/35576 (43) International Publication Date: 2 October 1997 (02.10.97)
(21) International Application Number: PCT/IL97/00085 (22) International Filing Date: 10 March 1997 (10.03.97) (30) Priority Data: 117702 28 March 1996 (28.03.96) IL (71) Applicant (for all designated States except US): TEL-AVIV MEDICAL CENTER RESEARCH AND DEVELOPMENT FUND [IL/IL]; 6 Weizman Street, 64239 Tel-Aviv (IL). (72) Inventors; and (75) Inventors/Applicants (for US only): ROTH, Arie [IL/IL]; 4 Efer Street, 69362 Tel-Aviv (IL). LANIADO, Shlomo [IL/IL]; 47 King David Boulevard, 64237 Tel-Aviv (IL). STERN, Naftali [IL/IL]; 17 Rimon Street, 72905 Nir-Zvi (IL). MAN, Abraham [IL/IL]; 4 Meir Yaqari Street, 69371 Tel-Aviv (IL). (74) Agent: REINHOLD COHN AND PARTNERS; P.O. Box 4060, 61040 Tel-Aviv (IL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>With amended claims and statement.</i>	
(54) Title: USE OF ISONIAZID FOR THE TREATMENT OF HYPERLIPOPROTEINEMIA			
(57) Abstract The known tuberculosis drug isoniazid (also known as isonicotinic acid hydrazide or INH) has unexpectedly been found to be useful for treating hyperlipoproteinemia. A method for decreasing the low-density lipoprotein/high-density lipoprotein (LDL/HDL) cholesterol ratio in the plasma of a mammal by administering isoniazid to the mammal is disclosed. Isoniazid can be administered together with other drugs known to lower the concentration of plasma lipoproteins to obtain an increased effect. A pharmaceutical composition containing isoniazid is also described.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF ISONIAZID FOR THE TREATMENT OF HYPERLIPOPROTEINEMIA

FIELD OF THE INVENTION

The present invention relates to the use of a known drug for the prevention and treatment of hyperlipoproteinemia which can lead to heart disease.

5

BACKGROUND OF THE INVENTION

Hyperlipoproteinemia is a condition in which the concentration of cholesterol- and/or triacylglycerol-carrying lipoproteins in the plasma exceeds a well defined normal limit. Clinical concern arises because an elevated concentration of lipoproteins can accelerate the development of atherosclerosis, thrombosis and infarction. Numerous population studies have shown that elevated serum concentration of total cholesterol, and especially low-density lipoprotein (LDL)-cholesterol, constitutes a major risk factor for the occurrence of atherosclerotic events. It has been clearly demonstrated that lowering the concentration of cholesterol-carrying lipoproteins in plasma can diminish the risk of myocardial infarction. However, there is now well established evidence that an increase in high-density lipoprotein (HDL)-cholesterol (and thus a decrease in the LDL/HDL cholesterol ratio) also has a beneficial and protective effect against the development of atherosclerosis and its complications. It is therefore more common to use the LDL/HDL cholesterol ratio as a measure of the plasma lipid profile with respect to predisposition to heart disease.

10
15
20

Various therapeutic measures are currently available in order to improve plasma lipid profile. These include the administration of various

drugs which generally lower the concentration of plasma lipoproteins (Goodman and Gilman. The Pharmacological Basis of Therapeutics, Eighth Edition (1990) Pergamon Press, Inc., Maxwell House, Fairview Park, Elmsford, New York, pg. 881-894) These drugs include:

- 5 (1) Fibrates: Aryloxyisobutyric acids and other related compounds collectively referred to as fibrates have been found to be effective in reducing plasma concentrations of triglycerides and LDL-cholesterol. Some of these drugs have also been shown to mildly increase HDL-cholesterol. Among the better known compounds are: clofibrate, gemfibrozil, fenofibrate, ciprofibrate, benzafibrate.
- 10 (2) HMG (hydroxymethylglutaric acid) CoA reductase inhibitors: These drugs, which are fungal-derived compounds, were found effective in lowering LDL-cholesterol and generally result in a mild increase in HDL-cholesterol. Compounds include: mevastatin, lovastatin, simvastatin, pravastatin.
- 15 (3) Nicotinic Acid: This drug decreases plasma triglycerides, LDL-cholesterol and induce a mild to moderate increase in HDL-cholesterol, but its use is limited by the frequent occurrence of serious side effects.

Most of the aforementioned drugs exert their beneficial effects mainly by reducing LDL-cholesterol. It would be advantageous to also have a drug which acts mainly to increase HDL-cholesterol without causing undue side effects.

20

Isoniazid, also known as isonicotinic acid hydrazide (INH), is considered to be a primary drug for the chemotherapy of tuberculosis (Goodman and Gilman, *supra*, pages 1146-1149). Other minor therapeutic uses which have been described for this drug include cerebellar tremor, Crohn's Disease, Huntington's Disease, Parkinson's Disease, Multiple Sclerosis and shoulder-hand syndrome. Although its precise anti-tuberculous mechanism is unknown, it was observed to act on the same hepatic enzymes as does alcohol. The latter has been shown to have the capability of increasing plasma HDL-cholesterol.

25

30

One study (Tanalp, R. and Uzalp, B., Ankara Univ. Tip Fak. Mecm. (1978), in Chem. Abst.91(1):243k) showed an increase in blood cholesterol in rats treated with a combination of isoniazid and testosterone propionate. Another study (Manyam, B.V. in Chest 84:120 (1983)) briefly described a decrease in serum cholesterol in humans on treatment with isoniazid. A
5 third study (Krishnamoorthy, M.S. and Karthikoyan, S. Pharmacol Res. (1991) 24(3):219-25) reported an increase in the levels of plasma total lipid and cholesterol in phenobarbitone sodium-pretreated rabbits after INH treatment, while treatment with INH alone did not alter the lipid levels in
10 plasma and tissues.

Thus, there is no clear indication in the literature as to a potential use of isoniazid in treating hyperlipoproteinemia.

BRIEF SUMMARY OF THE INVENTION

15 It is an object of the present invention to provide a drug capable of lowering the LDL/HDL plasma cholesterol ratio.

It is a further object of the present invention to provide a drug capable of increasing the HDL-cholesterol level in the plasma.

20 Additionally, it is an object of the present invention to provide a method of treating hyperlipoproteinemia in a mammal by drug administration.

It has now been surprisingly discovered that the anti-tuberculosis drug isoniazid is active in increasing the HDL cholesterol concentration in the plasma.

25 According to one aspect of the present invention, there is provided a method for decreasing the LDL/HDL cholesterol ratio in the plasma of a mammal comprising administering isoniazid to the mammal.

Further in accordance with this aspect of the present invention, there is provided a method for decreasing the LDL/HDL cholesterol ratio in the
30 plasma of a mammal wherein the ratio is decreased by an increase in the concentration of HDL-cholesterol in the plasma.

According to another aspect of the present invention, isoniazid is administered in combination with a hypolipidemic drug. The combination of the two drugs results in a lowering of the LDL/HDL ratio and enables utilizing lower doses of each of the individual drugs, thus minimizing drug specific side effects.

According to yet another aspect of the present invention, there is provided a use of isoniazid in the preparation of a pharmaceutical composition capable of decreasing the LDL/HDL cholesterol ratio in the plasma of a mammal.

DETAILED DESCRIPTION OF A PREFERRED EMBODIMENT

The effect of isoniazid in increasing the plasma HDL cholesterol concentration was demonstrated in the following clinical study. Nine apparently healthy individuals (six males and three females) were treated (because of suspected tuberculosis) with 300 mg/day of oral isoniazid for 6 months. Total cholesterol, HDL-cholesterol, LDL-cholesterol, and triglycerides were determined in serial blood tests taken at baseline, after 3 and 6 months of treatment, and 3 months after termination of the treatment (for which there are currently results for only 4 patients). The tests were carried out using kits (Boehringer Mannheim) based on an enzymatic hydrolysis method which measure triglycerides and cholesterol and HDL. Total cholesterol was calculated according to the Friedwald equation (Friedwald, Y. *et al.* Clin. Chem.. 18:499-502 (1972)).

The results are presented in Tables I-IV:

Table I: HDL-Cholesterol (mg/dl)

Patient	Age	Sex	Initial	3 months	6 months	3 month Post-treatment
1	69	F	69	71	69	...
2	58	F	40	44	50	42
3	22	M	56	54	59	...
4	17	F	62	97	98	65
5	58	M	53	44	62	52
6	50	M	68	72	92	...
7	73	M	40	39	54	...
8	38	M	57	74	71	...
9	43	M	47	52	62	49

Table II: LDL-Cholesterol (mg/dl)

Patient	Initial	3 months	6 months	3 months Post-treatment
1	103	113	105	...
2	132	125	155	42
3	76	62	92	...
4	82	84	90	98
5	144	123	152	162
6	80	95	109	...
7	134	110	120	...
8	151	121	117	...
9	99	88	95	129

Table III: Total Cholesterol (mg/dl)

Patient	Initial	3 months	6 months	3 month Post-treatment
1	209	201	195	...
2	228	225	260	234
3	145	125	164	...
4	159	194	199	179
5	226	191	240	240
6	162	194	231	...
7	229	188	207	...
8	245	212	191	...
9	183	163	201	196

Table IV: Triglycerides (mg/dl)

Patient	Initial	3 months	6 months	3 months Post-Treatment
1	184	158	165	...
2	280	281	275	428
3	66	47	64	...
4	74	66	54	79
5	144	121	128	131
6	71	133	151	...
7	276	194	161	...
8	185	85	64	...
9	161	116	95	129

The results are summarized in Table V:

Table V

parameter measured	baseline (n=9)	6 months (n=9)	p values
HDL-cholesterol	55±11	69±16	0.005
LDL-cholesterol	111±29	115±24	NS
total cholesterol	198±37	210±29	NS
triglycerides	160±82	129±70	0.165

NS - not significant

It may be seen that a significant increase in plasma HDL-cholesterol from 55 ± 11 mg% at baseline to 69 ± 16 mg% at 6 months ($p=0.005$) was attained. No significant changes were observed in the other parameters.

This clinical study was later broadened to encompass 15 patients. The HDL-cholesterol level was measured at the baseline and after 3 and 6 months of treatment. The results are summarized in Table VI:

Table VI: HDL-cholesterol (mg/dl)

months of treatment	baseline (n=15)	post-treatment (n=15)	p values
3	53 ± 11	61 ± 19	0.048
6	53 ± 11	65 ± 18	0.002

It can be seen that a significant increase in HDL-cholesterol was obtained with isoniazid treatment already after three months. Thus it can be seen that the administration of isoniazid results in a decrease in the LDL/HDL cholesterol ratio by increasing the HDL-cholesterol concentration.

The ratio can be further decreased by administering a hypolipidemic drug, which decreases the concentration of LDL cholesterol, in combination with isoniazid. Such drugs include fibric acids such as clofibrate, gemfibrozil, fenofibrate, ciprofibrate and bezafibrate; HMG CoA reductase inhibitors such as mevastatin, lovastatin, simvastatin and pravastatin; and nicotinic acid.

A pharmaceutical composition can be prepared for decreasing the LDL/HDL cholesterol ratio in the plasma of a mammal which comprises isoniazid together with a pharmaceutically acceptable excipient. Such excipients include for example amylan, acacia, tragacanth, sodium stearate glycolate, elcema, talc and calcium stearate. As discussed above, various hypolipidemic drugs can be included in the composition. The composition is preferably in an oral form. Typical doses of isoniazid are in the range of

- 10 -

10-300 mg accumulated dose. In the event that isoniazid is administered together with hypolipidemic drugs, the dose can be reduced.

5 The use of isoniazid in the treatment of hyperlipoproteinemia can provide a major contribution to the primary and secondary (post cardiovascular event) prevention of cardiovascular events, including myocardial infarction, cerebrovascular disease and peripheral vascular disease.

It will be appreciated by persons skilled in the art that the present invention is not limited to what has been thus far described, but rather the scope of the present invention is limited only by the following claims:

10

CLAIMS:

1. A method for decreasing the low-density lipoprotein/high-density lipoprotein (LDL/HDL) cholesterol ratio in the plasma of a mammal comprising administering isoniazid to said mammal.
- 5 2. A method according to claim 1 wherein said ratio is decreased by an increase in the concentration of HDL cholesterol in the plasma.
3. A method according to claim 2 wherein said ratio is further decreased by decreasing the concentration of LDL cholesterol in the plasma.
4. A method according to claim 1 comprising administering
10 isoniazid together with fibric acids.
5. A method according to claim 4 wherein said fibric acids are selected from the group consisting of clofibrate, gemfibrozil, fenofibrate, ciprofibrate and bezafibrate.
6. A method according to claim 1 comprising administering
15 isoniazid together with HMG CoA reductase inhibitors.
7. A method according to claim 6 wherein said HMG CoA reductase inhibitors are selected from the group consisting of mevastatin, lovastatin, simvastatin and pravastatin.
8. A method according to claim 1 comprising administering
20 isoniazid together with nicotinic acid to said mammal.
9. Isoniazid for use in the decrease of the LDL/HDL cholesterol ratio in the plasma of a mammal.
10. Isoniazid for use in the decrease of the LDL/HDL cholesterol ratio in the plasma of a mammal according to claim 9 whereby the ratio is
25 decreased by increasing the concentration of HDL cholesterol in the plasma.
11. Use of isoniazid in the preparation of a pharmaceutical composition for use in decreasing the LDL/HDL cholesterol ratio in the plasma of a mammal.
12. A use according to claim 11 wherein said ratio is decreased by
30 increasing the concentration of HDL cholesterol in the plasma.

- 12 -

13. A use according to claim 11 wherein said medicament further comprises fibric acids.
14. A use according to claim 11 wherein said medicament further comprises HMG CoA reductase inhibitors.
- 5 15. A use according to claim 11 wherein said medicament further comprises nicotinic acid.
16. A pharmaceutical composition for use in decreasing the LDL/-HDL cholesterol ratio in the plasma of a mammal comprising isoniazid and a pharmaceutically acceptable excipient.
- 10 17. A method for treating hyperlipoproteinemia in a mammal comprising administering isoniazid to said mammal.

AMENDED CLAIMS

[received by the International Bureau on 25 August 1997 (25.08.97);
original claims 1, 2, 9 - 12 and 16 amended;
remaining claims unchanged (2 pages)]

1. A method for increasing the concentration of HDL cholesterol in the plasma of a mammal comprising administering isoniazid to said mammal.
2. A method according to claim 1 wherein the low-density lipoprotein/high-density lipoprotein (LDL/HDL) cholesterol ratio in the plasma is decreased as a result of the increase in the concentration of HDL cholesterol in the plasma.
3. A method according to claim 2 wherein said ratio is further decreased by decreasing the concentration of LDL cholesterol in the plasma.
4. A method according to claim 1 comprising administering isoniazid together with fibric acids.
5. A method according to claim 4 wherein said fibric acids are selected from the group consisting of clofibrate, gemfibrozil, fenofibrate, ciprofibrate and bezafibrate.
6. A method according to claim 1 comprising administering isoniazid together with HMG CoA reductase inhibitors.
7. A method according to claim 6 wherein said HMG CoA reductase inhibitors are selected from the group consisting of mevastatin, lovastatin, simvastatin and pravastatin.
8. A method according to claim 1 comprising administering isoniazid together with nicotinic acid to said mammal.
9. Isoniazid for use in the increase of the concentration of HDL cholesterol in the plasma of a mammal.
10. Isoniazid for use in the increase of the concentration of HDL cholesterol according to claim 9 wherein the LDL/HDL cholesterol ratio is decreased by increasing the concentration of HDL cholesterol.
11. Use of isoniazid in the preparation of a pharmaceutical composition for use in increasing the concentration of HDL cholesterol in the plasma.

12. A use according to claim 11 wherein the LDL/HDL cholesterol ratio in the plasma of a mammal is decreased by increasing the concentration of HDL cholesterol in the plasma.
 13. A use according to claim 11 wherein said medicament further comprises fibric acids.
 14. A use according to claim 11 wherein said medicament further comprises HMG CoA reductase inhibitors.
 15. A use according to claim 11 wherein said medicament further comprises nicotinic acid.
 16. A pharmaceutical composition for use in increasing the concentration of HDL cholesterol in the plasma of a mammal comprising isoniazid and a pharmaceutically acceptable excipient.
 17. A method for treating hyperlipoproteinemia in a mammal comprising administering isoniazid to said mammal.
-

Statement under Article 19(1)

The amendment of the claims is based on making original Claim 2 the main claim and original Claim 1 a dependent claim. The claimed invention is based on the surprising discovery that isoniazid causes an increase in the concentration of HDL cholesterol in the plasma of a mammal. This discovery is not disclosed in any of the references cited in the Search Report.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/IL 97/00085

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/44 //(A61K31/44,31:215),(A61K31/44,31:19),(A61K31/44,31:35),(A61K31/44,31:22),(A61K31/44,31:44)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CAN. J. BIOCHEM. (1972), 50(1), 32-4 CODEN: CJBIAE, 1972, XP000654503 KUTTY, K. M. ET AL: "Serum cholinesterase activity in hyperlipidemia and the in vitro effect of isoniazid on serum cholinesterase"	1,3, 9-12,16, 17
Y	see the whole document	2,4-15
Y	DRUG CHEM. TOXICOL. (1977) (1991), 14(3), 293-303 CODEN: DCTODJ; ISSN: 0148-0545, 1991, XP002032797 KARTHIKEYAN, S. ET AL: "Effect of subacute administration of isoniazid and pyridoxine on lipids in plasma, liver and adipose tissues in the rabbit" see the whole document	2,9-12

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

24 June 1997

Date of mailing of the international search report

02.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrera, S

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IL 97/00085

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>JAMES E. E. REYNOLDS: "MARTINDALE THE EXTRA PHARMACOPOEIA" 1993, THE PHARMACEUTICAL PRESS, LONDON XP002032798 PAGES 984-994 AND 1045-1056 -----</p>	<p>4-8, 13-15</p>

THIS PAGE BLANK (USPTO)